Brain Research, 474 (1988) 407–409
Elsevier

Author Index

Akiguchi, I., see Yagi, H., 86
Alonso, M.J., see Fernández-Alfonso, M.S., 147

Ball, J., see Gibson, S.J., 255
Barnéoud, P., Neveu, P.J., Vitiello, S., Mormède, P. and Le Moal, M., Brain neocortex immunomodulation in rats, 394
Ben-Ari, Y., see Neuman, R.S., 201
Berman, K.F., see Iadarola, M.J., 75
Berry, M., see Sadler, M., 130
Björklund, A., see Kalén, P., 374
Blaustein, J.D., King, J.C., Toft, D.O. and Turcotte, J., Immunocytochemical localization of estrogen-induced progestin receptors in guinea pig brain, 1
Bloom, S.R., see Gibson, S.J., 255
Brownell, D.B., see Gibson, S.J., 255
Bush, K.T., see Nagele, R.G., 279

Cadet, J.L., see Iadarola, M.J., 75
Cau, P., see Pinard, R., 48
Cherubini, E., see Neuman, R.S., 201
Cronin, J. and Dudek, F.E., Chronic seizures and collateral sprouting of dentate mossy fibers after kainic acid treatment in rats, 181

Daddona, P.E., see Yamamoto, T., 112
De Belleroche, J., see Gibson, S.J., 255
De Biasi, S., see Spreafico, R., 192
Deckel, A.W., Moran, T.H. and Robinson, R.G., Receptor characteristics and recovery of function following kainic acid lesions and fetal transplants of the striatum: I. Cholinergic systems, 27
Deckel, A.W., Moran, T.H. and Robinson, R.G., Receptor characteristics and recovery of function following kainic acid lesions and fetal transplants of the striatum: II. Dopaminergic systems, 39
DeLanney, L.E., see Ricaurte, G.A., 359
Dewar, K., see Yamamoto, T., 112
Dow-Edwards, D.L., see Miller, M.W., 316
Dudek, F.E., see Cronin, J., 181
Duffy, L.K., see Joachim, C.L., 100

Endo, K., see Imai, H., 327

Fernández-Alfonso, M.S., Alonso, M.J., Rico, I., Salaices, M., Sánchez-Ferrer, C.F. and Marin, J., Effects of the Ca^{2+} agonists, Bay K 8644 and CGP 28392, on isolated cat cerebral and peripheral arteries, 147
Flores, C.M., see Iadarola, M.J., 75
Frassoni, C., see Spreafico, R., 192
Fuster, J.M., see Quintana, J., 211
Fuster, J.M., see Yajeya, J., 222

Gattaglia, G., see Spreafico, R., 192
Geiger, J.D., see Yamamoto, T., 112
Gibson, S.J., Polak, J.M., Katagiri, T., Su, H., Weller, R.O., Brownell, D.B., Holland, S., Hughes, J.T., Kikuyama, S., Ball, J., Bloom, S.R., Steiner, T.J., De Belleroche, J. and Rose, F.C., A comparison of the distributions of eight peptides in spinal cord from normal controls and cases of motor neurone disease with special reference to Onuf's nucleus, 255
Gillette, M.U. and Prosser, R.A., Circadian rhythm of the rat suprachiasmatic brain slice is rapidly reset by daytime application of cAMP analogs, 348
Gintzler, A.R., see Sander, H.W., 343
Goldstein, M., see Henschen, A.F., 66
Gondoh, Y., see Tsuda, A., 174
Gotoh, F., see Tomita, M., 165
Griffith, W.H. and Taylor, L., Sodium valproate decreases synaptic potentiation and epileptiform activity in hippocampus, 155

Hand, T.H., Koob, G.F., Stinus, L. and Le Moal, M., Aversive properties of opiate receptor blockade: evidence for exclusively central mediation in naive and morphine-dependent rats, 364
Harada, K., see Kato, I., 16
Harrell, L.E. and Parsons, D.S., The role of cortical sympathetic ingrowth in the behavioral effects of nucleus basalis magnocellularis lesions, 353
Hasegawa, T., see Kato, I., 16
Henschen, A.F., Goldstein, M. and Palmer, M.R., Evidence for functional contact between cografted locus coeruleus and spinal cord in oculo: electrophysiological studies, 66
Holland, S., see Gibson, S.J., 255
Hughes, J.T., see Gibson, S.J., 255
Hunter, E.T., see Nagele, R.G., 279

Iadarola, M.J., Flores, C.M., Cadet, J.L., Suddath, R., Illowsky, B. and Berman, K.F., Met^5-Enkephalin-Arg^6-Gly^7-Leu^8 immunoreactivity in rat and human cerebrospinal fluid: influence of neuroleptic drugs and electroconvulsive shock, 75
Ida, Y., see Tsuda, A., 174
Ikarashi, T., see Kato, I., 16
Illowsky, B., see Iadarola, M.J., 75
Imai, H., Nakamura, T., Endo, K. and Narabayashi, H., Hemiparkinsonism in monkeys after unilateral caudate nucleus infusion of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP): behavior and histology, 327
Imai, K., see Yamashita, T., 309
Irwin, I., see Ricaurte, G.A., 359
Ishida, M. and Shinozaki, H., Acromelic acid is a much more potent excitant than kainic acid or domoic acid in the isolated rat spinal cord, 386

Joachim, C.L., Duffy, L.K., Morris, J.H. and Selkoe, D.J., Protein chemical and immunocytochemical studies of meningovascular β-amyloid protein in Alzheimer's disease and normal aging, 100

Kalén, P., Kokaia, M., Lindvall, O. and Björklund, A., Basic characteristics of noradrenaline release in the hippocampus of intact and 6-hydroxydopamine-lesioned rats as studied by in vivo microdialysis, 374

Katagari, T., see Gibson, S.J., 255
Kato, I., Harada, K., Hasegawa, T. and Ikarashi, T., Role of the nucleus of the optic tract of monkeys in optokinetic nystagmus and optokinetic after-nystagmus, 16
Katoh, S., see Yagi, H., 86
Kawahara, K., Nakazono, Y., Kumagai, S., Yamauchi, Y. and Miyamoto, Y., Neuronal origin of parallel suppression of postural tone and respiration elicited by stimulation of mid-pontine dorsal tegmentum in the decerebrate cat, 403
Kikuyama, S., see Gibson, S.J., 255
King, J.C., see Blaustein, J.D., 1
Kitao, Y., see Tokuno, H., 390
Klaassen, A.B.M., see Van Megen, Y.J.B., 185
Kobari, M., see Tomita, M., 165
Kokaia, M., see Kalén, P., 374
Koob, G.F., see Hand, T.H., 364
Kosciuk, M.C., see Nagele, R.G., 279
Kovács, K.J. and Makara, G.B., Corticosterone and dexamethasone act at different brain sites to inhibit adrenalectomy-induced adrenocorticotropin hypersecretion, 205
Kudo, M., see Tokuno, H., 390
Kuijpers, W., see Van Megen, Y.J.B., 185
Kumagai, S., see Kawahara, K., 403
Kuraishi, Y., see Ohno, H., 197

Ladenheim, E.E., Speth, R.C. and Ritter, R.C., Reduction of CCK-8 binding in the nucleus of the solitary tract in unilaterally nodosectomized rats, 125
LaHoste, G.J., Mormède, P., Rivet, J.-M. and Le Moal, M., New evidence for distinct patterns of brain organization in rats differentiated on the basis of inherent laterality, 296
Langston, J.W., see Ricaurte, G.A., 359
Lanoir, J., see Pinard, R., 48
Lee, H.-y., see Nagele, R.G., 279
Le Moal, M., see Barnéoud, P., 394
Le Moal, M., see Hand, T.H., 364
Le Moal, M., see LaHoste, G.J., 296
Levy, S. and Tillotson, D., Effects of Na^+ and Ca^{2+} gradients on intracellular free Ca^{2+} in voltage-clamped *Aplysia* neurons, 333
Lindvall, O., see Kalén, P., 374

Makara, G.B., see Kovács, K.J., 205
Marin, J., see Fernández-Alfonso, M.S., 147
Miller, M.W. and Dow-Edwards, D.L., Structural and metabolic alterations in rat cerebral cortex induced by prenatal exposure to ethanol, 316
Minami, M., see Ohno, H., 197
Mitsuma, T., see Sobue, G., 287
Miyamoto, Y., see Kawahara, K., 403
Miyazaki, M., see Okuda, C., 399
Mizobe, T., see Okuda, C., 399
Moran, T.H., see Deckel, A.W., 27
Moran, T.H., see Deckel, A.W., 39
Moriizimi, T., see Tokuno, H., 390
Mormède, P., see Barnéoud, P., 394
Mormède, P., see LaHoste, G.J., 296
Morris, J.H., see Joachim, C.L., 100

Nagele, R.G., Kosciuk, M.C., Hunter, E.T., Bush, K.T. and Lee, H.-y., Immunoelectron microscopic localization of actin in neurites of cultured embryonic chick dorsal root ganglia: actin is a component of granular, microtubule-associated crossbridges, 279
Nagy, J.I., see Yamamoto, T., 112
Nakamura, T., see Imai, H., 327
Nakamura, Y., see Tokuno, H., 390
Nakazono, Y., see Kawahara, K., 403
Narabayashi, H., see Imai, H., 327
Neuman, R.S., Ben-Ari, Y. and Cherubini, E., Antagonism of spontaneous and evoked bursts by 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX) in the CA3 region of the in vitro hippocampus, 201
Neveu, P.J., see Barnéoud, P., 394
Nieminen, K., Suarez-Isla, B.A., Rapoport, S.I., Electrical properties of cultured dorsal root ganglion neurons from normal and trisomy 21 human fetal tissue, 246
Nordberg, A., see Wahlström, G., 189

Oguchi, M., see Tsuda, A., 174
Ohno, H., Kuraishi, Y., Minami, M. and Satoh, M., Modality-specific antinociception produced by intrathecal injection of anti-somatostatin antiserum in rats, 197
Okuda, C., Mizobe, T. and Miyazaki, M., Involvement of endogenous thyrotropin-releasing hormone in central regulation of the cardiovascular system after bleeding in conscious rats, 399

Palmer, M.R., see Henschen, A.F., 66
Pant, H.C., see Shah, J., 94
Parsons, D.S., see Harrell, L.E., 353
Pfeiffer, S.W., see Sawchenko, P.E., 231
Pinard, R., Segu, L., Cau, P. and Lanoir, J., Distribution of benzodiazepine receptors in the rat superior colliculus: a light and electron microscope quantitative autoradiographic study, 48
Polak, J.M., see Gibson, S.J., 255
Portoghese, P.S., see Sander, H.W., 343
Prosser, R.A., see Gillette, M.U., 348

Quintana, J., see Yajeya, J., 222
Quintana, J., Yajeya, J. and Fuster, J.M., Prefrontal representation of stimulus attributes during delay tasks. I. Unit activity in cross-temporal integration of sensory and sensory-motor information, 211

Rapoport, S.I., see Nieminen, K., 246
Ravid, R., see Van Zwieten, E.J., 369
Ricaurte, G.A., DeLanney, L.E., Wiener, S.G., Irwin, I. and Langston, J.W., 5-Hydroxyindoleacetic acid in cerebrospinal fluid reflects serotonergic damage induced by 3,4-methylenedioxymethamphetamine in CNS of non-human primates, 359
Rico, I., see Fernández-Alfonso, M.S., 147
Ritter, R.C., see Ladenheim, E.E., 125
Rivet, J.-M., see LaHoste, G.J., 296
Robinson, R.G., see Deckel, A.W., 27
Robinson, R.G., see Deckel, A.W., 39
Rodrigues de Miranda, J.F., see Van Megen, Y.J.B., 185
Rose, F.C., see Gibson, S.J., 255

Sadler, M. and Berry, M., Link-vertex analysis of Purkinje cell dendritic trees from the murine cerebellum, 130
Saito, N., see Yamashita, T., 309
Salaices, M., see Fernández-Alfonso, M.S., 147
Sánchez-Ferrer, C.F., see Fernández-Alfonso, M.S., 147
Sander, H.W., Portoghese, P.S. and Gintzler, A.R., Spinal κ-opiate receptor involvement in the analgesia of pregnancy:

effects of intrathecal nor-binaltorphimine, a κ-selective antagonist, 343
Satoh, M., see Ohno, H., 197
Sawchenko, P.E. and Pfeiffer, S.W., Ultrastructural localization of neuropeptide Y and galanin immunoreactivity in the paraventricular nucleus of the hypothalamus in the rat, 231
Segu, L., see Pinard, R., 48
Selkoe, D.J., see Joachim, C.L., 100
Shah, J. and Pant, H.C., Spontaneous calcium release induced by ethanol in the isolated rat brain microsomes, 94
Shinozaki, H., see Ishida, M., 386
Shirao, I., see Tsuda, A., 174
Sobue, G., Taki, T., Yasuda, T. and Mitsuma, T., Gangliosides modulate Schwann cell proliferation and morphology, 287
Speth, R.C., see Ladenheim, E.E., 125
Spreafico, R., de Biasi, S., Frassoni, C. and Gattaglia, G., A comparison of GAD- and GABA-immunoreactive neurons in the first somatosensory area (SI) of the rat cortex, 192
Staines, W.A., see Yamamoto, T., 112
Steiner, T.J., see Gibson, S.J., 255
Stinus, L., see Hand, T.H., 364
Su, H., see Gibson, S.J., 255
Suarez-Isla, B.A., see Nieminen, K., 246
Suddath, R., see Iadarola, M.J., 75
Swaab, D.F., see Van Zwieten, E.J., 369

Takeda, T., see Yagi, H., 86
Taki, T., see Sobue, G., 287
Tanaka, M., see Tsuda, A., 174
Taylor, L., see Griffith, W.H., 155
Tillotson, D., see Levy, S., 333
Toft, D.O., see Blaustein, J.D., 1
Tokuno, H., Moriizimi, T., Kudo, M., Kitao, Y. and Nakamura, Y., Ultrastructural differences between pallidohabenular terminals and pallidothalamic terminals, 390
Tomita, M., Gotoh, F. and Kobari, M., Colloid osmotic pressure of cat brain homogenate separated from autogenous CSF by a copper ferrocyanide membrane, 165
Tsuda, A., Tanaka, M., Ida, Y., Shirao, I., Gondoh, Y., Oguchi, M. and Yoshida, M., Expression of agression attenuates stress-induced increases in rat brain noradrenaline turnover, 174
Turcotte, J, see Blaustein, J.D., 1

Vaney, D.I. and Young, H.M., GABA-like immunoreactivity in NADPH-diaphorase amacrine cells of the rabbit retina, 380
Van Megen, Y.J.B., Klaassen, A.B.M., Rodrigues de Miranda, J.F. and Kuijpers, W., Cholinergic muscarinic receptors in rat cochlea, 185
Van Zwieten, E.J., Ravid, R., Swaab, D.F. and v.d. Woude, Tj., Immunocytochemically stained vasopressin binding sites on blood vessels in the rat brain, 369
Vitiello, S., see Barnéoud, P., 394

Wahlström, G. and Nordberg, A., Relation between voluntary ethanol intake in rats and changes in striatal muscarinic binding sites seen after induction of stable ethanol intake by an intermittent ethanol treatment, 189
Weller, R.O., see Gibson, S.J., 255
Wiener, S.G., see Ricaurte, G.A., 359
Woude, Tj. v.d., see Van Zwieten, E.J., 369

Yachi, A., see Yamashita, T, 309
Yagi, H., Katoh, S., Akiguchi, I. and Takeda, T., Age-related deterioration of ability of acquisition in memory and learning in senescence accelerated mouse: SAM-P/8 as an animal model of disturbances in recent memory, 86
Yajeya, J., Quintana, J. and Fuster, J.M., Prefrontal representation of stimulus attributes during delay tasks. II. The role of behavioral significance, 222
Yajeya, J., see Quintana, J., 211
Yamamoto, T., Staines, W.A., Dewar, K., Geiger, J.D., Daddona, P.E. and Nagy, J.I., Distinct adenosine deaminase-containing inputs to the substantia nigra from the striatum and tuberomammillary nucleus, 112
Yamashita, T., Imai, K., Saito, N. and Yachi, A., Detection of novel proteins associated with secondary amyloidosis and Alzheimer's disease by monoclonal antibody, 309
Yamauchi, Y., see Kawahara, K., 403
Yasuda, T., see Sobue, G., 287
Yoshida, M., see Tsuda, A., 174
Young, H.M., see Vaney, D.I., 380